arXiv:1204.6359v1 [astro-ph.CO] 28 Apr 2012

Hubble Constant, Lensing, and Time Delay in T*e*V*e* S

Yong Tian[1], Chung-Ming Ko[2], Mu-Chen Chiu[3]
[1]*Department of Physics, National Central University, Jhongli, Taiwan 320, R.O.C.*[*]
[2]*Institute of Astronomy, Department of Physics and Center for Complex Systems,*
National Central University, Jhongli, Taiwan 320, R.O.C.[†] *and*
[3]*Scottish University Physics Alliance, Institute for Astronomy, the Royal Observatory,*
University of Edinburgh, Blackford Hill, Edinburgh, EH9 3HJ, UK

(Dated: received 28 April 2012; published)

Hubble constant can be determined by the time delay of gravitational lensing systems. As data on time delay observation accumulates, it is time to revisit this approach. As in other dynamical phenomena in scales of galaxy and cluster of galaxies, gravitational lensing in these scales is also plagued by the problem of excess acceleration or gravity (a.k.a. missing mass problem). There are always some accelerations unaccounted for by luminous matter. Usually dark matter is introduced to interpret the discrepancy. However, MOdified Newtonian Dynamics (MOND) is more successful in explaining the excess accelerations in galaxy scale. We adopt T*e*V*e* S as the relativistic version of MOND to study gravitational lensing phenomena, and we can evaluate the Hubble constant from the derived time-delay formula. To apply our method, we rely on the CASTLE quasar lensing survey and the Sloan Digital Sky Survey (SDSS). Four samples are suitable for our study. Using only the luminous part of the lensing galaxies, the average of the derived Hubble constant is 68.5 km $\mathrm{s}^{-1}\mathrm{Mpc}^{-1}$.

Hubble constant H_0 is a long debated quantity in cosmology for more than half a century. Basically, it comes from the relation between cosmological distance and receding velocity of galaxies $v = H_0 d$. Its inverse represent the age of the universe.

The value of H_0 is sensitive to the way we estimate the distance. Its value has been estimated by many distance-determination methods, such as Cepheids, tip of the red giant branch, maser galaxies, surface brightness fluctuations, Tully-Fisher relation, Type Ia supernovae, and other methods such as gravitational time delay, Sunyaev-Zel'dovich (SZ) effect, cosmic microwave background (CMB), see ref [1, 2] for details. In this work, we focus on gravitational lensing and time delay. Obtaining H_0 by Gravitational time delay was introduced in 1964 by Refsdal [3]. Bright variable sources are needed and Refsdal suggested supernovae. Subsequent works on H_0, however, used quasar lensing. Now there are 19 systems of quasar lensing with time delay measurement [4, 5].

One advantage of using time delay to derive Hubble constant is that it is less sensitive to cosmological models. Thus it provides a more direct probe to cosmological distance [1]. However, there are some uncertainties in determining mass distribution by image deflections and distortions from gravitational lensing. This is commonly known as "mass sheet degeneracy" [6]. Another source of uncertainty in mass is, of course, the missing mass problem. Missing mass is a long standing issue. Oort in 1932 and Zwicky in 1933 were the first to put forward the notion of missing mass in galactic systems [7, 8]. The missing mass problem was neatly confirmed by the observed flat rotation curve in spiral galaxies [9–11]. Nowadays, missing mass exists in nearly all galactic systems, clusters of galaxies, large scale structure and CMB. In fact, the problem should be interpreted in terms of excess acceleration or gravity, i.e., there are some acceleration cannot be accounted for by luminous matter. To compensate the excess acceleration one can, on the one hand, introduce dark matter into the system. On the other hand, one can modify Newton's law of motion or law of gravity. In 1983, Milgrom proposed MOdified Newtonian Dynamics (MOND) to explain both flat rotation curve and Tully-Fisher relation [12]. MOND asserts that when acceleration is smaller than about $\mathfrak{a}_0 = 10^{-10}$ m s^{-2}, Newton's second law of motion has to be changed: $\tilde{\mu}(|\mathbf{a}|/\mathfrak{a}_0)\mathbf{a} = -\nabla\Phi_\mathrm{N} = \mathbf{a}_\mathrm{N}$, with $\tilde{\mu}(x) \approx x$ for $x \ll 1$, $\tilde{\mu}(x) \to 1$ for $x \gg 1$, where Φ_N is the Newtonian gravitational potential. $\tilde{\mu}(x)$ is called the interpolation function. For later calculation it is useful to introduce the inverted interpolation function $\tilde{\nu}$ such that $\mathbf{a} = -\nabla\Phi = \tilde{\nu}(|\mathbf{a}_\mathrm{N}|/\mathfrak{a}_0)\mathbf{a}_\mathrm{N}$

MOND is very successful in explaining the dynamics of galactic systems [13]. Recently, McGaugh showed that MOND can perfectly explain the Tully-Fisher relation in gas rich spiral galaxies without uncertainty parameter such as mass-to-light ratio [14, 15]. As usual the result created some debate [16–18]. Nevertheless, many consider MOND is not quite successful on cluster of galaxies scale [19]. A recent study on the gravitational redshift of clusters of galaxies [20] has generated some debate on whether MOND is applicable to cluster scale, and it seems that MOND does not have difficulty in interpreting the data [21]. In any case, the original MOND is a non-relativistic theory and can not be applied to relativistic phenomena such as gravitational lens and cosmology. Two decades after the original paper by Milgrom [12], in 2004 Bekenstein proposed the Tensor-Vector-Scalar (T*e*V*e* S) covariant relativistic gravity theory with MONDian dynamics as its non-relativistic limit [22]. In 2006, Chiu et al. started from T*e*V*e* S and derived the corresponding strong lens equation [23]. Later in 2009, Milgrom proposed another relativistic version of MOND called BiMOND [24]. It turns out that T*e*V*e* S and BiMOND have identical gravitational lensing results. The lensing theory has been applied to some galaxy lensing data, in which the galactic mass has been calculated and compared with population synthesis [25–27]. Once again, MOND can explain the data well. Moreover, the MONDian lensing mass is consistent with the dynamical mass from fundamental plane [28].

The observational tests above on galactic mass depend on the mass-to-light ratio, which cannot be measured directly. Here we put forward another test, namely, Hubble constant H_0 from time delay measurement. If MOND cannot get a reasonable value of H_0, it should be ruled out without any ambiguity. We solve the lens and time delay equation simultaneously to obtain both mass of the lens and the Hubble constant. Unlike dark matter theory with adjustable mass model, the mass distribution in MOND is fixed by the density profile of luminous matter.

The discussion on light deflection due to gravitational force can be traced to Newton's *Opticks*. Modern view of light deflection is a relativistic gravitational effect, in which both time-like and space-like part in the metric contribute to the deflection angle. Using General theory of Relativity (GR), Einstein derived the deflection angle which is just twice the Newtonian one. It turns out that the deflection angle derived from T*e*V*e* S or BiMOND is also twice the Newtonian one (i.e., the same as GR) except that the Newtonian potential is replaced by the MONDian potential. The deflection angle by a spherical lens in small angle approximation can be written as [23, 25]

$$\Delta\varphi = 2\int a_\perp \frac{dt}{c} \approx \frac{2\varrho_0}{c^2}\int_{-D_{\rm LS}}^{D'_{\rm L}} \frac{1}{\varrho}\frac{\partial\Phi(\varrho)}{\partial\varrho}\, d\zeta\,, \qquad (1)$$

where c is the speed of light, $\Phi(\varrho)$ is the MONDian potential, $\varrho_0 \approx D_{\rm L}\theta$ is the closest approach from the center of the lens, θ is the image position, and $\varrho^2 = \varrho_0^2 + \zeta^2$. $D_{\rm L}$, $D'_{\rm L}$ and $D_{\rm LS}$ are the angular distances of the lens from the observer, observer from the lens and the source from the lens, respectively. The direction of the image is in the direction of the closest approach (projected on the sky). In thin lens approximation the corresponding lensing equation is given by

$$\beta = \theta_+ - \alpha(\theta_+) = \alpha(\theta_-) - \theta_-\,, \qquad (2)$$

where β is the source position, $\theta_\pm$ are the two images on the same and opposite side as the source, respectively, and $\alpha(\theta) = \Delta\varphi D_{\rm LS}/D_{\rm S}$ is commonly called the reduced deflection angle.

Time delay is defined as the difference in time traveled by light along the actual path and along the undeflected path. It can be derived from Fermat principle or from the geodesic equation in relativistic gravitation theory. As in deflection angle, the form of the time delay is the same for GR and MOND (with Newtonian potential for GR and MONDian potential for MOND),

$$t(\theta) = \frac{(1+z_{\rm L})}{c}\left[\frac{D_{\rm L}D_{\rm S}}{2D_{\rm LS}}\alpha(\theta)^2 - \int_{-D_{\rm LS}}^{D'_{\rm L}} \frac{2\Phi(\varrho)}{c^2}\, d\zeta\right]\,. \qquad (3)$$

The first and second term in Eq. (3) are referred to the geometric and the potential time delay. The angular distance $D_{\rm A}$ is proportional to the Hubble constant, $D_{\rm A} = (c/H_0)\tilde{D}_{\rm A}(z)$, and $\tilde{D}_{\rm A}$ depends on redshift z only. In the cases where the difference in time delay of two images is available, the value of H_0 (and the mass of the lens as well) can be obtained by solving difference time delay equation $\Delta t = t(\theta_-) - t(\theta_+)$ and lens equation Eq. (2).

It is interesting to note that the time delay difference of MOND for point mass lens in Bekenstein form ($\boldsymbol{\nabla}\Phi = \boldsymbol{\nabla}\Phi_N(1+\sqrt{\mathfrak{a}_0/|\boldsymbol{\nabla}\Phi_N|})$) has exactly the same form as GR

$$\frac{\tilde{D}_{\rm LS}}{\tilde{D}_{\rm L}\tilde{D}_{\rm S}}\frac{H_0\Delta t}{(1+z_{\rm L})} = \frac{1}{2}\left(\theta_+^2 - \theta_-^2\right) + \theta_{\rm E}^2 \log\left(\frac{\theta_+}{\theta_-}\right)\,. \qquad (4)$$

Here $\theta_{\rm E} = \sqrt{4GMD_{\rm LS}/c^2 D_{\rm L}D_{\rm S}}$ is called the Einstein radius. The only difference is that lens equation gives $\theta_{\rm E}^2/\theta_+\theta_- = 1$ in GR, but in MOND it gives [23]

$$\frac{\theta_{\rm E}^2}{\theta_+\theta_-} + \frac{\pi}{(\theta_+ + \theta_-)}\frac{\theta_{\rm E}^2}{\theta_0} = 1\,, \qquad (5)$$

where $\theta_0 = \sqrt{GM/\mathfrak{a}_0 D_{\rm L}^2}$ (M is the mass of the luminous matter). We can use θ_0 as a criterion for deep MOND regime if the gravitational acceleration at the closest approach is much smaller than $\mathfrak{a}_0$, i.e., $\theta \gg \theta_0$. Because in Eq. (5), $\theta_{\rm E}^2 \to \theta_+\theta_-$ as $\theta_\pm \ll \theta_0$ (Newtonian) and $\theta_{\rm E}^2 \approx \theta_0(\theta_+ + \theta_-)/\pi$ as $\theta_\pm \gg \theta_0$ (MONDian) , from Eq. (4) the value of H_0 from MOND is smaller than that from GR, see Figure 1. In deep MOND regime the interpolation function is $\tilde{\mu}(a/\mathfrak{a}_0) \simeq a/\mathfrak{a}_0$. If the extend of the luminous matter is also much smaller than θ (i.e., practically a point mass), then the time delay difference is solely determined by the potential time delay, because the deflection angle approaches a constant in deep MOND regime [23]. The time delay difference is given by a simple formula which is independent of the choice of interpolation function,

$$\frac{\tilde{D}_{\rm LS}}{\tilde{D}_{\rm L}\tilde{D}_{\rm S}}\frac{H_0\Delta t}{(1+z_{\rm L})} = \frac{1}{2}\left(\theta_+^2 - \theta_-^2\right)\,. \qquad (6)$$

We emphasize that Eq. (6) do not have any free parameters for the interpolation function or mass of the lens, not even $\mathfrak{a}_0$ (the utmost important constant of MOND). $\mathfrak{a}_0$ remains in lensing Eq. (5) with mass of the lens only. Thus, if such a clean observation is available, then it has the potential to falsify MOND unambiguously.

In GR for Eq. (4), the first term is the geometric time delay and the second is the potential time delay. However, in MOND the first term in Eq. (4) comes from primary MONDian potential time delay $\pi\theta_{\rm E}^2(\theta_+ - \theta_-)/2\theta_0$. Becuase this primary MONDian contribution cancel the geometric time delay, the remain is the first term in (4). When we go from Newtonian regime ($\theta_0/\theta_\pm \gg 1$) to the deep MOND regime ($\theta_0/\theta_\pm \ll 1$), $\theta_{\rm E}$ changes from the order of θ to the order of $\sqrt{\theta_0\theta_\pm}$. The primary MONDian potential time delay remain finite, while the secondary MONDian potential time delay (the second term in Eq. (4)) diminish. As a result, in the deep MOND regime the time delay (Eq. (6)) becomes solely potential time delay.



FIG. 1. Hubble constant (km s^{-1} Mpc^{-1}) is plotted as a function of time delay difference (days) in a point mass lens model. The plot shows both the result of GR and deep MOND. Dashed line is GR, dotted-line is deep MOND in ΛCDM cosmology, solid line is deep MOND in massive neutrinos cosmology, and dot-dashed line for Hernquist model in MOND. To put the diagram into perspective, we consider a quasar lensing object FBQ J0951+2635. Its measured time delay difference is 16 days and is marked by the vertical black line. For reference $H_0 = 72$ km s^{-1} Mpc^{-1} is marked by the horizontal black line.

The gravitational lens with time delay measurement that we are able to collect are not in the deep MOND regime. The gravitational acceleration ranges from 10^{-9} ms^{-2} to 10^{-10} ms^{-2}. Thus, We need to pick an interpolation function. In literature several practical forms have been used, such as, Bekenstein form $\tilde{\mu}(x) = (-1 + \sqrt{1+4x})/(1+\sqrt{1+4x})$ [22], simple form $\tilde{\mu}(x) = x/(1+x)$ [29], and standard form $\tilde{\mu}(x) = x/\sqrt{1+x^2}$ [12]. Recently, a theoretical form from quantum effect in acceleration universe is proposed $\tilde{\mu}(x) = (\sqrt{4x^2+1}-1)/2x$ [30]. All these forms can be put into a canonical form [25]. The corresponding inverted canonical interpolation function is

$$\tilde{\nu}(x_{\rm N}) = \left[1 + \frac{1}{2}\left(\sqrt{4x_{\rm N}^{-\alpha} + \eta^2} + \eta\right)\right]^{1/\alpha}. \qquad (7)$$

In the following calculations we take the Bekenstein form, i.e., $(\alpha, \eta) = (1, 0)$.

TABLE I. Data of the four selected quasar lens with time delay measurement. The unit of $\theta_\pm$ is arsecond. H band effective radius [31] is used for HE 2149-2745 and FBQ J0951+2635, and V band for SDSS J0946+1835 [5]. Ellipticity $\epsilon = 1 - b/a$ are all measured in R band [5, 32, 33].

Name	z_l	z_s	θ''_+	θ''_-	θ''_e	ϵ	Δt(days)
HE 2149-2745	0.495	2.03	1.35	0.34	0.50	0.5	103±12
FBQ J0951+2635	0.240	1.25	0.88	0.22	0.17	0.25	16±2
SBS 0909+532	0.830	1.38	0.76	0.42	1.58	–	45^{+1}_{-11}
SDSS J0946+1835	0.388	4.81	2.43	0.64	2.59	0.16	142.7±6.8

Although hundreds of quasar lensing have been found, only a few have been measured with gravitational time delay. This method has been proven difficult because the amplitude of quasar variability is quite small, and clear and simple modeled relative image is hard to identify [1]. Up to now there are 19 strong lenses cases with measured time delay [4, 5]. In this paper, we consider spherical lens only. From the nineteen candidates, we select elliptical galaxy lensing systems with double images. Only 6 systems satisfy our criteria, they are HE 2149-275, FBQ J0951, SBS 0909+532, SDSS0946+1835 , SDSS J1650+4251,and HE 1104-1805. The rest includes clusters, spiral galaxies, multiple images or multiple lens, which require analysis beyond the scope of this paper and involve more uncertainty. Moreover, the lens galaxy in SDSS J1650+4251 is very dark. It does not have a reliable effective radius. And the uncertainty in time delay in HE 1104-1805 is too large. That brings our sample to a total of four systems. Table I lists some properties of the four selected systems.

The luminosity profile of the lens elliptical galaxies satisfies the de Vaucouleurs' profile. We, therefore, adopt the Hernquist mass density profile [34]. The Newtonian gravitational potential and acceleration of Hernquist model are $\Phi_{\rm N} = -GM/(\varrho + \varrho_{\rm h})$ and $\partial\Phi_{\rm N}/\partial\varrho = GM/(\varrho + \varrho_{\rm h})^2$. The Hernquist radius $\varrho_{\rm h}$ is 0.551 times the effective radius (half-light radius). The MONDian acceleration is $\partial\Phi/\partial\varrho = \tilde{\nu}(x_{\rm N})\partial\Phi_{\rm N}/\partial\varrho$ (where $x_{\rm N} = |\partial\Phi_{\rm N}/\partial\varrho|/\mathfrak{a}_0$).

Different cosmological models give different angular distance for the same redshift. As MOND is aiming at doing away with dark matter, thus, a cosmological model consistent with MOND is needed. MOND has been traditionally criticized that it can not form large-scale structure. Basically, the criticism originated from GR with baryons only. However, T*e*V*e* S contain additional fields which can form large-scale structure compatible with observations [35]. As for CMB, T*e*V*e* S needs 2eV massive neutrinos [36] which is treated as non-relativistic particle, $(\Omega_{\rm B}, \Omega_\nu, \Omega_\Lambda) = (0.05, 0.17, 0.78)$. We deem that Skordis et al. [36] gave a cosmological model consistent with MOND. The discrepancy between the two models is small in our sample. The difference in $\tilde{D}_{LS}/(\tilde{D}_L\tilde{D}_S)$ is within 0.4%.

TABLE II. The evaluated mass of the lens and Hubble constant. The unit of mass is 10^{10} solar mass and the unit of Hubble constant is km s^{-1} Mpc^{-1}. $x = a/\mathfrak{a}_0$ is a measure of the MONDian regime at the closest approach ϱ_0, and $x_\pm$ correspond to $\theta_\pm$.

	Mass		H_0			
Name	GR	MOND	GR	MOND	x_-	x_+
HE 2149-2745	$23.2^{25.8}_{20.5}$	$16.1^{17.5}_{14.6}$	$72.2^{81.7}_{64.7}$	$57.6^{66.0}_{51.0}$	11.3	2.3
FBQ J0951+263	$2.9^{3.3}_{2.6}$	$2.3^{2.5}_{2.1}$	$93.2^{106.5}_{82.9}$	$79.3^{91.5}_{69.8}$	28.5	4.0
SBS 0909+532	$77.2^{78.9}_{58.3}$	$55.8^{56.8}_{44.3}$	$84.5^{111.9}_{82.7}$	$70.2^{95.0}_{68.5}$	8.7	5.9
SDSS J0946+1835	$95.3^{99.8}_{90.7}$	$64.1^{66.4}_{61.8}$	$84.9^{89.1}_{81.0}$	$67.0^{70.7}_{63.6}$	7.8	2.8

In this treatment, we don't have any free parameter. Only mass and Hubble constant are unknown variable to be solved. Flat rotation curve of spiral galaxies and Tully-Fisher relation in gas rich galaxies give a consistent value of the acceleration constant $\mathfrak{a}_0 = 1.21 \times 10^{-10}$ m s^{-2} [13, 14, 29]. As we select lensing galaxies that satisfy Hernquist's model and the Hernquist radius would be determined by photometric measurement of the effective radius, the only unknown in the mass model is the total galactic mass. On the part of gravitational lensing, the angular position and the redshift (of the lens and the two images) and the time delay difference between the two images are measured. The only unknown left in the lensing part is the Hubble constant. We can evaluate two unknowns by solving the lens equation and time delay equation simultaneously. The result is summarized in Table II. The smaller numbers in the first four columns are results taken into account of corresponding upper and lower uncertainties.

$x = a/\mathfrak{a}_0$ in the last two columns of Table II is the ratio of the acceleration at the closest approach to the MOND acceleration constant. Recall that $x \gg 1$ is the Newtonian regime and $x \ll 1$ is the deep MOND regime. On the one hand, $x \lesssim 1$, MOND gives less mass than GR. The excess mass in GR can be interpreted as missing mass or dark matter. On the other hand, if $x \gg 1$, MOND and GR should give similar mass. As shown in Table II, most x is of the order of unity (intermediate MOND regime). As expected the mass and Hubble constant from MOND are smaller than the value computed in GR without dark matter. Once again, as in other galactic scale dynamical phenomena, MOND gives a consistent picture of explaining the observed excess acceleration in gravitational lensing including time delay phenomenon.

The other source of uncertainty comes from the choice of interpolation function. In Newtonian regime $x \gg 1$ or the deep MOND regime $x \ll 1$, different interpolation functions should give the same result. However, our sample lies in the intermediate MOND regime. Table II shows the result from Bekenstein form. Other interpolation functions would give somewhat different result. In fact, major uncertainty comes from observation, in particular, the time delay measurements.

In summary, this work is a first attempt to use MOND to interpret data from gravitational time delay. The Hubble constant evaluated from this study is consistent with literature (see Table II). A simple average of these four cases give $H_0 = 68.5$ km $s^{-1}Mpc^{-1}$. When compare with GR (without dark matter), the evaluated mass of the lens is 28% to 49% smaller than those from GR, and the Hubble constant is 18% to 27% smaller than GR. We stress that the ellipticity of HE 2149-2745 is too large to consider it as a spherical potential. We should interpret its result with caution. Among the four systems we have studied, the data from SDSS0946+1835 is the best as it has the least uncertainty in time delay, which is about 5%. The uncertainty it contributes to mass is about 3.7% and Hubble constant about 5%. Perhaps the longer the time delay, the smaller the measurement uncertainty will be. We look forward to more data from lens with long time delay. Gravitational lensing promises to provide a testing ground for modified gravity.

We thank Ting-Hung Peng for his help in data collection. This work is supported in part by the Taiwan National Science Council Grants NSC 98-2923-M-008-01-MY3 and NSC 99-2112-M-008-015-MY3.

---

[*] yonngtian@gmail.com

[†] cmko@astro.ncu.edu.tw

[1] W.L. Freedman, & B.F. Madore, Annu. Rev. Astron. Astrophys. **48**, 673 (2010)

[2] N. Jackson, Living Rev. Relativity, 10, (2007), 4; http://www.livingreviews.org/lrr-2007-4

[3] S. Refsdal, Mon. Not. Roy. Astron. Soc. **128**, 307 (1964)

[4] P. Danuta, & H.Jens, Astrophys. J. **712**, 1378 (2010)

[5] I.D. McGreer, et al., Astrophys. J. **140**, 370 (2010)

[6] M.V. Gorenstein, I.I. Shapiro, & E.E. Falco, Astrophys. J. **327**, 693 (1988)

[7] J.H. Oort, Bulletin of the Astronomical Institutes of the Netherlands **6**, 249 (1932)

[8] F. Zwicky, Helv. Phys. Acta **6**, 110 (1933)

[9] V. Rubin and W.K. Fort, Astrophys. J. **159**, 379 (1970)

[10] T.S. Van Albada, J.N. Bahcall, K. Begeman, & R. Sanscisi, Astrophys. J. **295**, 305 (1985)

[11] K.G. Begman, A&A **223**, 47 (1989)

[12] M. Milgrom, Astrophys. J. **270**, 365 (1983)

[13] R.H. Sanders, & S.S. McGaugh, Ann. Rev. Astron. Astrophys. **40**, 263 (2002)

[14] S.S. McGaugh, Phys. Rev. Lett. **106**, 121303 (2011)

[15] S.S. McGaugh, Astron. J. **140**, 40 (2012)

[16] S. Foreman, & D. Scott, Phys. Rev. Lett. **108**, 141302 (2012)

[17] N.Y. Gnedin, arXiv:1108.2271v1[astro-ph.CO] (2011)

[18] S.S. McGaugh, arXiv:1109.1599v1[astro-ph.CO] (2011)

[19] A. Aguirre, J. Schaye, E. Quataert, Astrophys. J. **561**, 550 (2001)

[20] R. Wojtak, S.H. Hansen, & J. Hjorth, Nature **477**, 567 (2011)

[21] J.D. Bekenstein, & R.H. Sanders, Mon. Not. Roy. Astron. Soc **421**, L59 (2012)

[22] J.D. Bekenstein, Phys. Rev. D **70**, 083509 (2004)

[23] M.C. Chiu, C.M. Ko, & Y. Tian, Astrophys. J. **636**, 565 (2006)

[24] M. Milgrom, Phys. Rev. D **80**, 123536 (2009)

[25] M.C. Chiu, C.M. Ko, Y. Tian, & H.S Zhao, Phys. Rev. D **83**, 063523 (2011)

[26] H.S. Zhao, D.J. Bacon, A.N. Taylor, & K. Horne, Month. Not. Roy. Astron. Soc. **368**, 171 (2006)

[27] I. Ferreras, M. Sakellariadou, & M.F. Yusaf, Phys. Rev. Lett. **100**, 031302 (2008)

[28] R.H. Sanders, & D.D. Land, Mon. Not. Roy. Astron. Soc. **389**, 701 (2008)

[29] B. Famaey, & J. Binney, Mon. Not. R. Astron. Soc. **363**, 603 (2005)

[30] C.M. Ho, D. Minic, & Y.J. Ng, Physics Letters B **693**, 567 (2010)

[31] C.S. Kochanek, et al., Astrophys. J. **543**, 131 (2000)

[32] S. Lopez, O. Wucknitz, and L. Wisotzki, A&A **339**, 13 (1998)

[33] P. Jakobsson, et al., A&A **431**, 103 (2005)

[34] L. Hernquist, Astrophys. J. **356**, 359 (1990)

[35] C. Skordis, Class. Quantum Grav. **26**, 143001 (2009)

[36] C. Skordis, et al., Phys. Rev. Lett. bf96, 011301 (2006)